arXiv:1412.7700v2 [math.GR] 25 Feb 2015

TORSION SUBGROUPS IN THE UNITS OF THE INTEGRAL GROUP RING OF $\mathrm{PSL}(2,p^3)$

ANDREAS BÄCHLE AND LEO MARGOLIS

Abstract. We show that for every prime r all r-subgroups in the normalized units of the integral group ring of $\mathrm{PSL}(2,p^3)$ are isomorphic to subgroups of $\mathrm{PSL}(2,p^3)$. This answers a question of M. Hertweck, C.R. Höfert and W. Kimmerle for this series of groups.

Let G be a finite group and $\mathbb{Z}G$ its integral group ring. A natural question is how far the structure of G is reflected in the structure of the torsion part of $\mathrm{V}(\mathbb{Z}G)$, the group of normalized units of the integral group ring of G, and vice versa. One of the first questions studied in this context is, whether a finite subgroup of $\mathrm{V}(\mathbb{Z}G)$ is necessarily isomorphic to a subgroup of G, which was already raised in G. Higman's PhD thesis [Hig40, Section 5] (cf. also the article [San81] outlining this thesis). It was later explicitly stated as a problem in [San85, Problem 5.4].

A first step towards an answer to the above question lies in the consideration of p-subgroups. It is known that if G has cyclic Sylow p-subgroups then all p-subgroups of $\mathrm{V}(\mathbb{Z}G)$ are cyclic; for $p=2$ see [Kim07, Proposition] (the proof involves the Brauer-Suzuki theorem) and cf. [Her08, Corollary 1] for odd primes p. If the Sylow 2-subgroups of G are elementary abelian, then so are all 2-subgroups of $\mathrm{V}(\mathbb{Z}G)$ and in particular they are isomorphic to subgroups of G (combining [CL65, Corollary 4.1] and [Sak71, Corollary 1.7], cf. also [Seh93, Lemma (37.3)]). But it is not even known in general whether abelian Sylow p-subgroups of G cause that the p-subgroups of $\mathrm{V}(\mathbb{Z}G)$ are abelian. However, this assertion holds for solvable G [DJ96, Proposition 2.11].

In the concluding remarks of [HHK09] the question whether the p-subgroups of $\mathrm{V}(\mathbb{Z}\,\mathrm{PSL}(2,p^f))$ are (elementary) abelian is highlighted. For $p=2$ or $f\leq 2$ there is an even stronger assertion, recently proved by the second author [Mar14]: for all primes r the r-subgroups of $\mathrm{V}(\mathbb{Z}\,\mathrm{PSL}(2,p^f))$ are conjugate by a unit of the corresponding rational group algebra to a subgroup of the group base $\mathrm{PSL}(2,p^f)$. It is also known that any subgroup of $\mathrm{V}(\mathbb{Z}\,\mathrm{PSL}(2,p^f))$ of the same order as $\mathrm{PSL}(2,p^f)$ is conjugate in the rational group algebra to the group base $\mathrm{PSL}(2,p^f)$ [Ble95, Propositions 3.2 and 4.1].

In this article we prove that for all primes p the p-subgroups of the group $\mathrm{V}(\mathbb{Z}\,\mathrm{PSL}(2,p^3))$ are abelian. We obtain the following result:

Theorem. *Let p be a prime and set $G=\mathrm{PSL}(2,p^3)$. Then for all primes r, the r-subgroups of $\mathrm{V}(\mathbb{Z}G)$ are isomorphic to subgroups of G.*

The method of choice here is what one could call a "non-cyclic HeLP-method". For a finite group G consider a finite subgroup U of $\mathrm{V}(\mathbb{Z}G)$. Every ordinary representation of G can be linearly extended to a representation of $\mathbb{Z}G$ and then restricted to a representation D of $\mathrm{V}(\mathbb{Z}G)$. Let its character be denoted by χ. If ψ is a character of U, then for the inner product we necessarily have

$$\langle \chi|_U, \psi\rangle_U \in \mathbb{Z}_{\geq 0}.$$

For cyclic U this can be expressed in a explicit formula, whose application is known as the HeLP-method, cf. [LP89, Theorem 1] and [Her07, Section 4]. The method for non-cyclic U was used for the first time in the PhD-thesis of C.R. Höfert [Höf08, page 58] and later in [Her08] and [HHK09, page 2].

Date: February 26, 2015.

2010 *Mathematics Subject Classification.* 16U60, 20C05 (primary) and 16S34 (secondary).

Key words and phrases. integral group ring, torsion units, projective special linear group, p-subgroups.

The first author is supported by the Research Foundation Flanders (FWO - Vlaanderen).

We will introduce some notation. Let $u = \sum_{g\in G} z_g g \in \mathbb{Z}G$ be a normalized torsion unit. For x^G, the conjugacy class of the element $x \in G$ in G, we denote by

$$\varepsilon_x(u) = \sum_{g\in x^G} z_g$$

the *partial augmentation of u at x*. The partial augmentations will provide us with restrictions on the possible eigenvalues of $D(u)$ and vice versa. By the so-called Berman-Higman Theorem $\varepsilon_1(u) = 0$ if $u \neq 1$ [Hig40, Theorem 10], [Ber55, Lemma 2] (cf. also [Seh93, Proposition (1.4)]). Also $\varepsilon_x(u) = 0$ if the order of x does not divide the order of u [Her07, Proposition 2.2]. Moreover, the order of a finite subgroup of $\mathrm{V}(\mathbb{Z}G)$ divides the order of G [Sak71, Corollary 1.7], cf. also [Seh93, Lemma (37.3)], and its exponent divides the exponent of G [CL65, Corollary 4.1]. We will use these facts in the sequel without further mention.

Let A be a complex matrix of finite order. Assume that A has eigenvalues $\alpha_1, ..., \alpha_j$ each with multiplicity m_1, eigenvalues $\beta_1, ..., \beta_k$ each with multiplicity m_2 and eigenvalues $\gamma_1, ..., \gamma_\ell$ each with multiplicity m_3, then we indicate this by

$$A \sim \left(m_1 \times \boxed{\alpha_1, ..., \alpha_j}, m_2 \times \boxed{\beta_1, ..., \beta_k}, m_3 \times \boxed{\gamma_1, ..., \gamma_\ell}\right).$$

Proof of the theorem.

Set $G = \mathrm{PSL}(2, p^3)$. By a result of Dickson [Dic58, 260.] (see also [Hup67, 8.27 Hauptsatz]), the Sylow r-subgroups of the simple groups $\mathrm{PSL}(2, p^f)$ are elementary abelian for $r = p$, cyclic for odd $r \neq p$, and dihedral groups if $r = 2 \neq p$. By the results cited above we obtain that the r-subgroups of $\mathrm{V}(\mathbb{Z}G)$ are isomorphic to subgroups of G, provided the Sylow r-subgroups are cyclic. The case of elementary abelian Sylow 2-subgroups is also handled by the remarks in the introduction. If the Sylow 2-subgroups are dihedral, the result is obtained in [HHK09, Theorem 2.1]. It remains the case $r = p \geq 3$. Note that in this case the Sylow p-subgroups of G are elementary abelian of order p^3.

Let H be a finite p-subgroup of $\mathrm{V}(\mathbb{Z}G)$. Hence $|H| \leq p^3$ and $\exp H \mid p$. Assume that H is not isomorphic to a subgroup of G, then, by the classification of all p-groups up to order p^3, it is a so-called Heisenberg group. Thus there are elements $z, b, c \in H$ such that

$$H = \langle z,\ b,\ c \mid z^p = b^p = c^p = 1,\ z \in \mathrm{Z}(H),\ c^{-1}bc = zb\rangle \simeq (C_p \times C_p) \rtimes C_p. \tag{1}$$

We will use the non-cyclic HeLP-method to show that H does not exist. In G there are exactly two conjugacy classes of elements of order p, let g and h be representatives of these classes. In Table 1 we list two irreducible characters [Dor71, Theorem 38.1], one of them we will use in the remainder of the proof. Let $\epsilon \in \{1, -1\}$ such that $p \equiv \epsilon \bmod 4$.

	1	g	h
η	$\frac{p^3+\epsilon}{2}$	$\frac{\epsilon+\sqrt{\epsilon p^3}}{2}$	$\frac{\epsilon-\sqrt{\epsilon p^3}}{2}$
η'	$\frac{p^3+\epsilon}{2}$	$\frac{\epsilon-\sqrt{\epsilon p^3}}{2}$	$\frac{\epsilon+\sqrt{\epsilon p^3}}{2}$

Table 1. Part of the character table of $G = \mathrm{PSL}(2, p^3)$, $p \geq 3$, $\epsilon \in \{1, -1\}$ such that $p \equiv \epsilon \bmod 4$.

Let $\zeta = \exp(2\pi i/p) \in \mathbb{C}$, a primitive p-th root of unity, Q be a set of integral representatives of the quadratic residues in $(\mathbb{Z}/p\mathbb{Z})^\times$ and N be a set of integral representatives of the non-quadratic residues in $(\mathbb{Z}/p\mathbb{Z})^\times$. We will also use the Gaussian sums cf. [Gau11] (also [Neu99, Proof of (8.6) in Chapter I])

$$\begin{aligned} \sqrt{\epsilon p} &= 1 + 2\sum_{q\in Q} \zeta^q, \\ -\sqrt{\epsilon p} &= 1 + 2\sum_{n\in N} \zeta^n. \end{aligned} \tag{2}$$

Let D be a representation affording η. Let $u \in H \setminus \{1\}$ and $\alpha \in \mathbb{Z}_{\geq 0}$ such that $(\varepsilon_g(u), \varepsilon_h(u)) = (\alpha+1, -\alpha)$ or $(\varepsilon_g(u), \varepsilon_h(u)) = (-\alpha, \alpha+1)$. By slight abuse of notation we denote from here on by η also the restriction $\eta|_H$. For the first possibility of the partial augmentations of u we obtain, using the first equation of (2),

$$\begin{aligned}\eta(u) &= (\alpha+1)\left(\frac{\epsilon+\sqrt{\epsilon p^3}}{2}\right) + (-\alpha)\left(\frac{\epsilon-\sqrt{\epsilon p^3}}{2}\right)\\ &= \frac{1}{2}\left(\epsilon + (2\alpha+1)p\sqrt{\epsilon p}\right)\\ &= \frac{p+\epsilon}{2} + \alpha p + (2\alpha+1)p\sum_{q\in Q}\zeta^q. \end{aligned} \tag{3}$$

For the second possibility for the partial augmentations of u we get, using the second equation of (2),

$$\eta(u) = \frac{p+\epsilon}{2} + \alpha p + (2\alpha+1)p\sum_{n\in N}\zeta^n. \tag{4}$$

From the character value we get $\frac{p+\epsilon}{2} + \alpha p + \left(\frac{p-1}{2}\right)(2\alpha+1)p$ eigenvalues of $D(u)$. The other eigenvalues have to sum up to 0, hence there are

$$\begin{aligned}&\frac{1}{p}\left(\frac{p^3+\epsilon}{2} - \left(\frac{p+\epsilon}{2} + \alpha p + \left(\frac{p-1}{2}\right)(2\alpha+1)p\right)\right)\\ =\ &\frac{1}{2p}\left(p^3+\epsilon-p-\epsilon-2\alpha p-(p-1)(2\alpha+1)p\right)\\ =\ &\frac{p^2-p}{2} - \alpha p\end{aligned}$$

blocks having eigenvalues $1, \zeta, \zeta^2, ..., \zeta^{p-1}$. Thus in the case of $\varepsilon_g(u) > 0$ we have

$$D(u) \sim \left(\left(\frac{p+\epsilon}{2}+\alpha p\right)\times\boxed{1}, (2\alpha+1)p\times\boxed{\zeta^{q_1}, ..., \zeta^{q_{\frac{p-1}{2}}}}, \left(\frac{p^2-p}{2}-\alpha p\right)\times\boxed{1, \zeta, ..., \zeta^{p-1}}\right)$$

where $Q = \left\{q_1, ..., q_{\frac{p-1}{2}}\right\}$. In the other case, $\varepsilon_g(u) \leq 0$, we get

$$D(u) \sim \left(\left(\frac{p+\epsilon}{2}+\alpha p\right)\times\boxed{1}, (2\alpha+1)p\times\boxed{\zeta^{n_1}, ..., \zeta^{n_{\frac{p-1}{2}}}}, \left(\frac{p^2-p}{2}-\alpha p\right)\times\boxed{1, \zeta, ..., \zeta^{p-1}}\right)$$

with $N = \left\{n_1, ..., n_{\frac{p-1}{2}}\right\}$.

The group H is an extra-special p-group, its character theory is well-known, see e.g. [Dor71, Theorem 31.5]. H has exactly $p-1$ non-linear irreducible characters which are all of degree p. They all vanish on the non-central elements of H and take the values $p\zeta^j$ for $1 \leq j \leq p-1$ on the non-trivial central elements. Moreover H posses p^2 linear characters corresponding to the quotient $H/\mathrm{Z}(H) = H/H' \simeq C_p \times C_p$. In particular, they have value 1 on all central elements of H. See table 2. Now we decompose $\eta = \eta|_H$ into the irreducible characters of H.

Let $v \in \mathrm{Z}(H) \setminus \{1\}$. To obtain the multiplicity of the eigenvalue 1 of $D(v)$ as calculated above, we must sum up exactly

$$\frac{p+\epsilon}{2} + \alpha p + \frac{p^2-p}{2} - \alpha p = \frac{p^2+\epsilon}{2}$$

linear characters of H and hence

$$\frac{1}{p}\left(\frac{p^3+\epsilon}{2} - \frac{p^2+\epsilon}{2}\right) = \frac{p^2-p}{2}$$

irreducible non-linear character of H.

Now let $w \in H$ be a non-central element. Since every irreducible non-linear character of H vanishes on w, the character value $\eta(w)$ is the sum of exactly $\frac{p^2+\epsilon}{2}$ roots of unity. Thus from (3)

	1	b	b^2	...	$b^{p-1}c^{p-1}$	z	...	z^{p-1}
χ_1						1	...	1
χ_2						1	...	1
$\vdots$			$\mathrm{CT}(C_p \times C_p)$			$\vdots$		$\vdots$
χ_{p^2-1}						1	...	1
χ_{p^2}						1	...	1
ψ_1	p	0	0	...	0	$p\zeta$	...	$p\zeta^{p-1}$
$\vdots$	$\vdots$				$\vdots$	$\vdots$		$\vdots$
ψ_{p-1}	p	0	0	...	0	$p\zeta^{p-1}$	...	$p\zeta$

TABLE 2. Character table of $H \simeq (C_p \times C_p) \rtimes C_p$, the box indicates the character table of the quotient group $H/\mathrm{Z}(H) \simeq C_p \times C_p$.

and (4) we obtain that $(\varepsilon_g(w), \varepsilon_h(w)) = (1,0)$ or $(\varepsilon_g(w), \varepsilon_h(w)) = (0,1)$. Furthermore, from the eigenvalues of $D(u)$ calculated in these equations, we get

$$(\varepsilon_g(w), \varepsilon_h(w)) = (1,0) \qquad \Leftrightarrow \qquad (\varepsilon_g(w^n), \varepsilon_h(w^n)) = (0,1) \quad \forall n \in N. \tag{5}$$

Now we compute the inner product $\langle \eta, \chi \rangle_H$ of η with a non-trivial linear character χ of H. We split up the computation of the contributions of different parts of H and omit the global factor $1/|H| = 1/p^3$ until adding all contributions. The contribution of the identity element is

$$\eta(1)\chi(1) = \frac{p^3 + \epsilon}{2}.$$

There are $p-1$ more elements in the center of H. On η half of them takes the value $\frac{p+\epsilon}{2} + \alpha p + (2\alpha+1)p \sum_{q \in Q} \zeta^q$, the other half the value $\frac{p+\epsilon}{2} + \alpha p + (2\alpha+1)p \sum_{n \in N} \zeta^n$. Since they all lie in the kernel of χ the contribution to the inner product is

$$\begin{aligned} &\frac{p-1}{2}\left(\frac{p+\epsilon}{2} + \alpha p + (2\alpha+1)p \sum_{q \in Q} \zeta^q\right) + \frac{p-1}{2}\left(\frac{p+\epsilon}{2} + \alpha p + (2\alpha+1)p \sum_{n \in N} \zeta^n\right) \\ &= \frac{p-1}{2}\Big(p + \epsilon + 2\alpha p - (2\alpha+1)p\Big) \\ &= \epsilon \cdot \frac{p-1}{2}. \end{aligned}$$

The kernel of χ contains another conjugacy class of cyclic subgroups consisting of exactly p subgroups. In every such cyclic subgroup $\frac{p-1}{2}$ of the elements take the value $\frac{p+\epsilon}{2} + p \sum_{q \in Q} \zeta^q$ on η and $\frac{p-1}{2}$ of the elements take the value $\frac{p+\epsilon}{2} + p \sum_{n \in N} \zeta^n$ on η. The same computation as above gives the contribution

$$p\epsilon \cdot \frac{p-1}{2}.$$

To calculate the contributions of the other elements we need the following formulas which are direct consequences of the Gaussian sums in (2):

$$\left(\sum_{q \in Q} \zeta^q\right)\left(\sum_{i \in Q} \zeta^i\right) + \left(\sum_{n \in N} \zeta^n\right)\left(\sum_{j \in N} \zeta^j\right) = \frac{\epsilon \cdot p + 1}{2}$$

and

$$\left(\sum_{n \in N} \zeta^n\right)\left(\sum_{i \in Q} \zeta^i\right) + \left(\sum_{q \in Q} \zeta^q\right)\left(\sum_{j \in N} \zeta^j\right) = \frac{-\epsilon \cdot p + 1}{2}.$$

Every conjugacy class of non-central cyclic subgroups contains exactly p subgroups. Let $\langle d\rangle$ be such a subgroup. By $\chi(d)\in Q$ we indicate that $\chi(d)=\zeta^q$ for some $q\in Q$. To compute the contribution of the remaining elements we distinguish two cases.

Case 1: $(\varepsilon_g(d),\varepsilon_h(d))=(1,0)$, $\chi(d^{-1})\in Q$ or $(\varepsilon_g(d),\varepsilon_h(d))=(0,1)$, $\chi(d^{-1})\in N$. Then the contribution of the conjugacy class of $\langle d\rangle$ is

$$\begin{aligned}
p\sum_{h\in\langle d\rangle\setminus\{1\}}\eta(h)\chi(h^{-1}) &= p\left(\sum_{q\in Q}\left(\frac{p+\epsilon}{2}+p\sum_{i\in Q}\zeta^i\right)\zeta^q+\sum_{n\in N}\left(\frac{p+\epsilon}{2}+p\sum_{j\in N}\zeta^j\right)\zeta^n\right)\\
&= p\left(-\frac{p+\epsilon}{2}+p\left(\left(\sum_{q\in Q}\zeta^q\right)\left(\sum_{i\in Q}\zeta^i\right)+\left(\sum_{n\in N}\zeta^n\right)\left(\sum_{j\in N}\zeta^j\right)\right)\right)\\
&= p\left(-\frac{p+\epsilon}{2}+p\left(\frac{\epsilon\cdot p+1}{2}\right)\right)\\
&= \epsilon\cdot\frac{p^3-p}{2}
\end{aligned}$$

Case 2: $(\varepsilon_g(d),\varepsilon_h(d))=(1,0)$, $\chi(d^{-1})\in N$ or $(\varepsilon_g(d),\varepsilon_h(d))=(0,1)$, $\chi(d^{-1})\in Q$. Then the contribution of the conjugacy class of $\langle d\rangle$ is

$$\begin{aligned}
&p\left(\sum_{n\in N}\left(\frac{p+\epsilon}{2}+p\sum_{i\in Q}\zeta^i\right)\zeta^n+\sum_{q\in Q}\left(\frac{p+\epsilon}{2}+p\sum_{j\in N}\zeta^j\right)\zeta^q\right)\\
&= p\left(-\frac{p+\epsilon}{2}+p\left(\left(\sum_{n\in N}\zeta^n\right)\left(\sum_{i\in Q}\zeta^i\right)+\left(\sum_{q\in Q}\zeta^q\right)\left(\sum_{j\in N}\zeta^j\right)\right)\right)\\
&= p\left(-\frac{p+\epsilon}{2}+p\left(\frac{-\epsilon\cdot p+1}{2}\right)\right)\\
&= -\epsilon\cdot\frac{p^3+p}{2}
\end{aligned}$$

From now on let $I=\{0,1,...,p-1\}$. Let χ be a non-trivial linear character of H and let $s\in\mathrm{Ker}(\chi)\setminus\mathrm{Z}(H)$. Moreover let $t\notin\mathrm{Ker}(\chi)$ such that $\chi(t^{-1})\in Q$. Then $\{(ts^i)^{-1}\mid i\in I\}$ is a set which contains exactly one element from every conjugacy class of cyclic subgroups not lying in $\mathrm{Ker}(\chi)$ and $\chi((ts^i)^{-1})\in Q$ for every i. Set

$$\gamma=|\{\ i\mid(\varepsilon_g(ts^i),\varepsilon_h(ts^i))=(1,0)\ \}|$$

and

$$\delta=|\{\ i\mid(\varepsilon_g(ts^i),\varepsilon_h(ts^i))=(0,1)\ \}|.$$

Then $\gamma+\delta=p$ and, summing up all the contributions obtained above, we get

$$\begin{aligned}
\langle\eta,\chi\rangle_H &= \frac{1}{p^3}\left(\frac{p^3+\epsilon}{2}+\epsilon\cdot\frac{p-1}{2}+\epsilon p\cdot\frac{p-1}{2}+\epsilon\gamma\cdot\frac{p^3-p}{2}-\epsilon\delta\cdot\frac{p^3+p}{2}\right)\\
&= \frac{1}{2p^3}\left(p^3\left(1+\epsilon\gamma-\epsilon\delta\right)+\epsilon p^2-\epsilon(\gamma+\delta)p\right)\\
&= \frac{1+\epsilon\gamma-\epsilon\delta}{2}.
\end{aligned}$$

For $n\in N$ the map $\chi^n\colon H\to\mathbb{Z}[\zeta]\colon x\mapsto\chi(x)^n$ is also a linear character of H and an analogous computation gives

$$\langle\eta,\chi^n\rangle_H=\frac{1-\epsilon\gamma+\epsilon\delta}{2}.$$

Since both, $\langle\eta,\chi\rangle_H$ and $\langle\eta,\chi^n\rangle_H$, are non-negative, necessarily $|\gamma-\delta|\leq 1$. Thus $\gamma=\frac{p\pm1}{2}$ and $\delta=\frac{p\mp1}{2}$.

Recall that by (1), H is generated by a central element z and two other elements b and c. We may assume w.l.o.g. by (5) that $(\varepsilon_g(c),\varepsilon_h(c))=(1,0)$. We have that $\{\langle c\rangle,\langle bc^i\rangle\mid i\in I\}$ is a set

of representatives of the $p+1$ conjugacy classes of non-central cyclic subgroups of H. Up to the action of $\mathrm{Gal}(\mathbb{Q}(\zeta)/\mathbb{Q})$, every linear character is determined by its kernel. Let $n \in N$. Then by (5) there exist $a_0, a_1, ..., a_{p-1} \in \{1, n\}$ such that $c, b^{a_0}, b^{a_1}c^{a_1}, ..., b^{a_{p-1}}c^{(p-1)a_{p-1}}$ all have partial augmentation 1 at g and 0 at h.

Let χ be a non-trivial linear character with $\langle c\rangle \subseteq \mathrm{Ker}(\chi)$ and $\chi(b^{-1}) \in Q$. Then $S = \{bc^i \mid i \in I\}$ is a set which contains exactly one element from every conjugacy class of cyclic subgroups not lying in $\mathrm{Ker}(\chi)$. By the above, $\frac{p\pm1}{2}$ of the elements of S have partial augmentation 1 at g and the other elements have augmentation 0 at g. Since $\varepsilon_g(bc^i) = 1$ if and only if $a_i = 1$ we get

$$|\{\ i \in I \mid a_i = 1\ \}| \in \left\{\frac{p\pm1}{2}\right\}. \tag{6}$$

Now let $j \in I$ and χ be a non-trivial linear character of H such that $\langle bc^j\rangle \subseteq \mathrm{Ker}(\chi)$ and $\chi(c^{-1}) \in Q$. For every $i \in I \setminus \{j\}$ we determine one element of the form $b^{\ell a_i}c^{\ell i a_i}$, for some ℓ, lying in the coset $\mathrm{Ker}(\chi)c = \left\{z^r b^k c^{jk+1} \mid 0 \le k, r \le p-1\right\}$. To do so let ℓ, k be such that

$$b^k c^{jk+1} = b^{\ell a_i}c^{\ell i a_i}.$$

Thus $k \equiv \ell a_i \bmod p$ and $jk + 1 \equiv \ell i a_i \bmod p$. This gives $\ell \equiv (ia_i - ja_i)^{-1} \bmod p$. For the partial augmentations of $b^{\ell a_i}c^{\ell i a_i}$ it only matters, by (5), whether ℓ is a quadratic residue modulo p. Hence $b^{\ell a_i}c^{\ell i a_i}$ has the same partial augmentations as c if and only if $\ell \in Q$, i.e. $ia_i - ja_i \in Q$. Hence

$$1 + |\{\ i \in I \setminus \{j\} \mid ia_i - ja_i \in Q\ \}| \in \left\{\frac{p\pm1}{2}\right\}, \tag{7}$$

where the 1 represents the element c.

Denote by $(\ r \mid p\ )$ the Legendre symbol of r modulo the prime p and set $\beta_i = (\ a_i \mid p\ )$ for $i \in I$. By (6) we have

$$\sum_{i\in I}\beta_i = \sum_{i\in I}(\ a_i \mid p\ ) \in \{\pm1\}. \tag{8}$$

From (7) we get for every $j \in I$:

$$1 + \sum_{i\in I}(\ i-j \mid p\ )\beta_i = 1 + \sum_{i\in I}(\ (i-j)a_i \mid p\ ) \in \{\pm1\}.$$

Set $s_i = (\ i \mid p\ )$ for $i \in \{1, ..., p-1\}$. Then the above equations read as follows: there exist $m_0, ..., m_{p-1} \in \{\pm1\}$ such that

$$\begin{pmatrix} \beta_0 & s_1 & s_2 & \dots & s_{p-1} \\ s_{p-1} & \beta_1 & s_1 & \dots & s_{p-2} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ s_2 & s_3 & \dots & \beta_{p-2} & s_1 \\ s_1 & s_2 & \dots & s_{p-1} & \beta_{p-1} \end{pmatrix} \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_{p-2} \\ \beta_{p-1} \end{pmatrix} = \begin{pmatrix} m_0 \\ m_1 \\ \vdots \\ m_{p-2} \\ m_{p-1} \end{pmatrix}.$$

Summing up all equations and sorting with regard to the β_i's we get

$$\beta_0(s_1 + ... + s_{p-1}) + ... + \beta_{p-1}(s_1 + ... + s_{p-1}) + p = m_0 + ... + m_{p-1}.$$

Since $s_1 + ... + s_{p-1} = 0$ all the m_i's must be equal to 1. Thus we have to solve the homogeneous system of linear equations given by

$$\begin{pmatrix} 0 & s_1 & s_2 & \dots & s_{p-1} \\ s_{p-1} & 0 & s_1 & \dots & s_{p-2} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ s_2 & s_3 & \dots & 0 & s_1 \\ s_1 & s_2 & \dots & s_{p-1} & 0 \end{pmatrix} \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_{p-2} \\ \beta_{p-1} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}.$$

The null space of the system has (at most) dimension 1 by the following lemma and contains the vector $(\beta_0, \beta_1, ..., \beta_{p-1})^t = (1, 1, ..., 1)^t$. This is however out of the question by (8). □

Lemma. *Let m be odd and $A \in \mathbb{F}_2^{m\times m}$ where*

$$A = \begin{pmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & 0 & 1 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 1 & \dots & 0 & 1 \\ 1 & 1 & \dots & 1 & 0 \end{pmatrix}.$$

Then $\operatorname{rank} A = m - 1$.

Proof. Summing up all rows of A gives the zero vector, hence the rank of A is at most $m-1$. Let $z_1, ..., z_{m-1}$ be the first $m-1$ rows of A and $\alpha_1, ..., \alpha_{m-1} \in \mathbb{F}_2$ such that $\sum_{i=1}^{m-1} \alpha_i z_i = 0$. Then we get $\sum_{i=1, i\neq j}^{m-1} \alpha_i = 0$ for every $j \in \{1, ..., m-1\}$ and $\sum_{i=1}^{m-1} \alpha_i = 0$. Hence $\alpha_j = 0$ for all $j \in \{1, ..., m-1\}$. □

References

[Ber55] S.D. Berman, *On the equation $x^m = 1$ in an integral group ring*, Ukrain. Mat. Ž. **7** (1955), 253–261.

[Ble95] F.M. Bleher, *Tensor products and a conjecture of Zassenhaus*, Arch. Math. (Basel) **64** (1995), no. 4, 289–298.

[CL65] J.A. Cohn and D. Livingstone, *On the structure of group algebras I*, Canadian Journal of Mathematics **17** (1965), 583–593.

[Dic58] L.E. Dickson, *Linear groups: With an exposition of the Galois field theory*, with an introduction by W. Magnus, Dover Publications, Inc., New York, 1958.

[DJ96] M.A. Dokuchaev and S.O. Juriaans, *Finite subgroups in integral group rings*, Canad. J. Math. **48** (1996), no. 6, 1170–1179.

[Dor71] L. Dornhoff, *Group representation theory. Part A: Ordinary representation theory*, Marcel Dekker, Inc., New York, 1971, Pure and Applied Mathematics, 7.

[Gau11] C.F. Gauß, *Summatio quarumdam serierum singularium*, Commentationes soc. reg. sc. Gotting. recentiores **1** (1811), Published in German in: Gauss, Untersuchungen über höhere Arithmethik, Chelsea Publishing Co., New York, 1965.

[Her07] M. Hertweck, *Partial Augmentations and Brauer character values of torsion units in group rings*, `arXiv:math.RA/0612429v2[math.RA]` (2007).

[Her08] ———, *Unit groups of integral finite group rings with no noncyclic abelian finite p-subgroups*, Comm. Algebra **36** (2008), no. 9, 3224–3229.

[HHK09] M. Hertweck, C.R. Höfert, and W. Kimmerle, *Finite groups of units and their composition factors in the integral group rings of the group* PSL$(2, q)$, J. Group Theory **12** (2009), no. 6, 873–882.

[Hig40] G. Higman, *Units in group rings*, D. phil. thesis, Oxford Univ., 1940.

[Höf08] C.R. Höfert, *Bestimmung von Kompositionsfaktoren endlicher Gruppen aus Burnsideringen und ganzzahligen Gruppenringen*, Doktorarbeit, Universität Stuttgart, 2008, `http://elib.uni-stuttgart.de/opus/frontdoor.php?source_opus=3541&la=de`.

[Hup67] B. Huppert, *Endliche Gruppen I*, Die Grundlehren der mathematischen Wissenschaften ed., vol. 134, Springer-Verlag, Berlin, 1967.

[Kim07] W. Kimmerle, *Torsion units in integral group rings of finite insoluble groups*, Oberwolfach Reports **4** (2007), no. 4, 3229–3230, Abstracts from the mini-workshop held November 25–December 1, 2007, Organized by Eric Jespers, Zbigniew Marciniak, Gabriele Nebe, and Wolfgang Kimmerle.

[LP89] I.S. Luthar and I.B.S. Passi, *Zassenhaus conjecture for A_5*, Proc. Indian Acad. Sci. Math. Sci. **99** (1989), no. 1, 1–5.

[Mar14] L. Margolis, *A Sylow theorem for the integral group ring of PSL(2,q)*, `arXiv:1408.6075[math.RA]` (2014).

[Neu99] J. Neukirch, *Algebraic number theory*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 322, Springer-Verlag, Berlin, 1999.

[Sak71] A. I. Saksonov, *Group rings of finite groups. I*, Publ. Math. Debrecen **18** (1971), 187–209.

[San81] R. Sandling, *Graham Higman's thesis "Units in group rings"*, Integral representations and applications (Oberwolfach, 1980), Lecture Notes in Math., vol. 882, Springer, Berlin-New York, 1981, pp. 93–116.

[San85] ———, *The isomorphism problem for group rings: a survey*, Orders and their applications (Oberwolfach, 1984), Lecture Notes in Math., vol. 1142, Springer, Berlin, 1985, pp. 256–288.

[Seh93] S.K. Sehgal, *Units in integral group rings*, Pitman Monographs and Surveys in Pure and Applied Mathematics, vol. 69, Longman Scientific & Technical, Harlow, 1993.

Vakgroep Wiskunde, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussels, Belgium
E-mail address: `abachle@vub.ac.be`

Fachbereich Mathematik, Universität Stuttgart, Pfaffenwaldring 57, 70569 Stuttgart, Germany
E-mail address: `leo.margolis@mathematik.uni-stuttgart.de`